

Form for the Diagnosis of Death using Neurological Criteria in Children >2 months to <18 years {full guidance version}

This form is consistent with and should be used in conjunction with, the AoMRC (2008) A Code of Practice for the Diagnosis and Confirmation of Death¹ and has been endorsed for use by the following institutions: Faculty of Intensive Care Medicine, Intensive Care Society, Paediatric Intensive Care Society and National Organ Donation Committee: Paediatric Subgroup.

Number HOSPITAL ADDRESSOGRAPH or
Surname
First Name
Date of Birth
NHS

Objective of Care

- To diagnose and confirm the death of a mechanically ventilated, severely brain injured patient in coma, using neurological criteria.

Academy of the Medical Royal Colleges Definition of Human Death (2008)¹

“Death entails the irreversible loss of those essential characteristics which are necessary to the existence of a living human person and, thus, the definition of death should be regarded as the irreversible loss of the capacity for consciousness, combined with irreversible loss of the capacity to breathe. The irreversible cessation of brain-stem function whether induced by intra-cranial events or the result of extra-cranial phenomena, such as hypoxia, will produce this clinical state and therefore irreversible cessation of the integrative function of the brain-stem equates with the death of the individual and allows the medical practitioner to diagnose death.”

Context

- National professional guidance advocates the confirmation of death by neurological criteria wherever this seems a likely diagnosis and regardless of the likelihood of organ donation.^{3,4}
- UK General Medical Council (GMC) guidance on end of life care (2010) states that national procedures for identifying potential organ donors should be followed and, in appropriate cases, the specialist nurse for organ donation (SN-OD) should be notified. NICE guidance and UKPICS Standards recommend that the specialist nurse for organ donation (SN-OD) should be notified at the point when the clinical team declare the intention to perform brain-stem death tests.^{2,3,4}

Date and time of referral to SN-OD:

- Whilst most patients will already be in an Intensive Care Unit (ICU) when the diagnosis is suspected, some patients may be in other areas, e.g. the Emergency Department. On such occasions, it is legitimate, if considered necessary, to transfer a patient to the ICU for the diagnosis to be made.
- For many clinicians, the diagnosis and confirmation of death using neurological criteria, will be a relatively infrequent task and may be complicated by uncertainties regarding the nature of the primary diagnosis, irreversibility and the availability of suitably experienced personnel. Updated guidance on the diagnosis and confirmation of death by neurological criteria was published by the Academy of the Medical Royal Colleges (AoMRC) in 2008. ¹ For infants from 37 weeks to <2 months guidance was published by the Royal College of Paediatric and Child Health in 2015. ⁶

The person with parental responsibility for the child and other close family members should be made aware that the purpose of testing is to confirm if the child’s death has already occurred. If given an opportunity to witness the neurological examination, they should be appropriately supported and prepared for the possibility of spinal reflexes and their relevance, as far as the diagnosis of death by neurological criteria is concerned. Whether the child’s close family witness the clinical examination or not, the child’s need for dignity, privacy and spiritual support, should remain paramount.

Form for the Diagnosis of Death using Neurological Criteria in Children >2 months to <18 years {full guidance version}

Patient Name:

NHS Number:

Preparation		
<p>1. Evidence for Irreversible Brain Damage of known Aetiology Case records, past medical history including possibly contacting the GP, relevant imaging.</p>		
<p>2. Exclusion of Reversible Causes of Coma and Apnoea Standard ICU cardio-respiratory monitoring (to ensure haemodynamic stability), medication chart and history, blood and urine drug assay results (where relevant), drug antagonists (e.g. flumazenil, naloxone), peripheral nerve stimulator, recent serum glucose and biochemistry, thermometer, patient warming device.</p>		
<p>3. Tests for Absence of Brain-Stem Function <i>Brain-stem reflexes:</i> Bright light source; small sterile gauze swabs, otoscope with disposable ear pieces, 20 or 50 ml luer lock syringe and disposable quill, ice-cold water; a spatula, Yankauer sucker or laryngoscope, endotracheal suction catheters. <i>Apnoea test:</i> Haemodynamic monitoring (continuous ECG, invasive arterial pressure), arterial blood gas analysis including blood gas syringes x4, pulse oximetry and end-tidal CO₂ monitoring, means of delivering oxygen to the trachea by bulk flow (i.e. using a circuit which maintains oxygenation and CPAP e.g. mapleson C, ayres T piece, or endotracheal suction catheter and oxygen tubing).</p>		
<p>Diagnostic caution is advised in the following 'Red Flag' patient groups. (Based on the literature and unpublished case reports.) For advice in difficult circumstances contact the local or regional Clinical Lead for Organ Donation or the regional paediatric intensive care units.</p>		
<p>1. Testing <6 hours of the loss of the last brain-stem reflex</p>	<p>4. Patients with any neuro-muscular disorders</p>	<p>6. Prolonged fantanyl infusions</p>
<p>2. Testing <24 hours where aetiology primarily anoxic damage</p>	<p>5. Steroids given in space occupying lesions such as abscesses</p>	<p>7. Aetiology primarily located to the brain-stem or posterior fossa</p>
<p>3. Hypothermia (24-hour observation period following re-warming to normothermia recommended)</p>		

Examining Doctors	
Date and time:	Patient Location
Doctor One, Name and Designation	Doctor Two, Name and Designation
Name:	Name:
Grade:	Grade:
<p>Guidance</p> <p>1. The diagnosis of death by neurological criteria should be made by at least two medical practitioners. Both medical practitioners should have been registered with the General Medical Council (or equivalent Professional Body) for more than five years and be competent in the assessment of a patient who may be deceased following the irreversible cessation of brain-stem function and competent in the conduct and interpretation of the brain-stem examination. Both doctors should be competent in the diagnosis of death by neurological criteria. In infants, children and young adolescents it is recommended that one of the doctors should be a paediatrician and at least one should be a consultant.</p> <p>2. Those carrying out the tests must not have, or be perceived to have, any clinical conflict of interest and neither doctor should be a member of the transplant team. Clinical Leads for Organ Donation can carry out testing and are likely to have significant expertise.</p> <p>3. Testing should be undertaken by the nominated doctors acting together and must always be performed on two occasions. A complete set of tests should be performed on each occasion, i.e., a total of two sets of tests will be performed. Doctor One may perform the tests while Doctor Two observes; this would constitute the first set. Roles may be reversed for the second set. The tests, in particular the apnoea test, are therefore performed only twice in total.</p>	

Form for the Diagnosis of Death using Neurological Criteria in Children >2 months to <18 years {full guidance version}

Patient Name:

NHS Number:

Evidence for Irreversible Brain Damage of known Aetiology
<p>Diagnosis:</p> <p>Evidence for Irreversible Brain Damage of known Aetiology:</p> <p>Guidance</p> <ol style="list-style-type: none"> 1. The patient must have a Glasgow Coma Score of 3 and be mechanically ventilated with apnoea. 2. There should be no doubt that the patient's condition is due to irreversible brain damage of known aetiology. 3. It remains the duty of the two doctors carrying out the testing to be satisfied with the aetiology, the exclusion of all potentially reversible causes, the clinical tests of brain-stem function and of any ancillary investigations (see page 8); so that each doctor may independently confirm death following irreversible cessation of brain-stem function. 4. Occasionally it may take a period of continued clinical observation and investigation to be confident of the irreversible nature of the brain injury. The timing of the first test and the timing between the two tests should be adequate for the reassurance of all those directly concerned. 5. It is recommended that there is a minimum of twenty-four hours, of continued clinical observation, in patients where anoxic damage, following cardiorespiratory arrest, is the aetiology of the brain injury. If prior treatment of the patient has included induced hypothermia, it is recommended that there is a minimum of twenty-four hours, of continued clinical observation, following re-warming to normothermia. See above for 'Red Flag' patient groups. 6. Stabilisation of the patient prior to testing, especially support of the cardiovascular system, is a prerequisite to testing. Mean blood pressure should be consistently maintained at age appropriate levels (see appendix 1)⁵ and appropriate fluid resuscitation administered. This almost invariably requires the use of inotropes / vasopressors via central venous access. 7. Diabetes insipidus can develop rapidly and should be suspected in patients with a high urine output (typically greater than 4 ml/kg/hr) and rising Na⁺. Matched urinary and plasma electrolytes and osmolality may assist in the diagnosis. Treatment with desmopressin, 2 month- 12 years 400 nanograms, 12 to 18 years 1 to 4 micrograms IV boluses, is usually sufficient for treatment but repeated doses or vasopressin infusion may be required. Serum sodium should ideally be maintained between 140-160 mmol/L.

Exclusion of Reversible Causes of Coma and Apnoea		
<p>Guidance</p> <p>It is necessary to maintain circulation and respiration in the preceding hours prior to testing, with the aim of achieving relatively normal cardiovascular and respiratory physiological parameters. ⁵</p> <p>Suggested cardiovascular and respiratory goals for children:</p> <ul style="list-style-type: none"> • Sinus rhythm 60-170 beats per minute, as age appropriate (see appendix 1)⁵ • Mean arterial pressure 44-111 mmHg, as age appropriate (see appendix 1)⁵ • Central Venous Pressure 4 – 10 mmHg • Cardiac Index >2.5 – 6 l/min/m² (if measured) • Mixed venous oxygen saturation >60% - 70 % (if measured) • Maintenance of normocarbia and avoidance of hypoxia, acidaemia or alkalaemia. 		
	1st Test	2nd Test
<p>Mean arterial pressure at time of testing? Should be consistently maintained at age appropriate levels prior to testing.</p>	mmHg	mmHg

Form for the Diagnosis of Death using Neurological Criteria in Children >2 months to <18 years {full guidance version}

Patient Name:

NHS Number:

PaCO₂ at time of testing? A goal of normocarbica (PaCO ₂ <6.0 kPa), <i>if possible</i> , is recommended in the preceding hours prior to testing. See below for <i>starting PaCO₂ in the apnoea test</i> .	kPa	kPa		
PaO₂ at time of testing? Hypoxia should be avoided (PaO ₂ >10 kPa).	kPa	kPa		
Arterial pH/[H⁺] at time of testing? Acidaemia and alkalaemia should be avoided, <i>if possible</i> , aiming for a relatively normal pH 7.35-7.45 / [H ⁺] 45-35 nmol/L.	pH/[H ⁺]=	pH/[H ⁺]=		
Is the coma or apnoea due to ongoing cardio-respiratory instability? (To diagnose death using neurological criteria, ALL answers should be NO)	Dr One	Dr Two	Dr One	Dr Two

Guidance

The patient should not be receiving any drugs that might be contributing to the unconsciousness, apnoea and loss of brain-stem reflexes (narcotics, hypnotics, sedatives or tranquillisers); nor should they have any residual effect from any neuromuscular blocking agents (atracurium, vecuronium or suxamethonium).

It remains the duty of the two doctors carrying out the testing to be satisfied that sufficient time has elapsed to ensure that any remaining drug effect is non-contributory to the unconsciousness and loss of brain-stem reflexes. This will be based on an assessment of the medications the patient has received and from knowledge of the pharmacokinetics of these agents. Renal or hepatic failure and immaturity in younger children may prolong metabolism / excretion of these drugs. See above for 'Red Flag' patient groups.

	1 st Test		2 nd Test	
Where there is any doubt, specific drug levels should be measured (midazolam should be less than <10mcg/L, thiopentone <5mg/L).	Drug levels (if measured):		Drug levels (if measured):	
Antagonists such as flumazenil, naloxone and neostigmine may be used but there is no specific pharmacological data for predicting the dose effect of these antagonists.	Drug antagonists (if used)		Drug antagonists (if used)	
Residual neuromuscular blockade can be tested for, if felt necessary, by peripheral nerve stimulation.	Train of Four (if measured):		Train of Four (if measured):	
Is the coma or apnoea due to depressant drugs? (To diagnose death using neurological criteria, ALL answers should be NO)	Dr One	Dr Two	Dr One	Dr Two
Body temperature at time of testing? If core temperature is ≤ 34°C testing cannot be carried out.	°C		°C	

Form for the Diagnosis of Death using Neurological Criteria in Children >2 months to <18 years {full guidance version}

Patient Name:

NHS Number:

Serum sodium (Na⁺) at time of testing? Serum sodium should be between 115-160mmol/L. Rapid rises or falls in Na ⁺ should be avoided	mmol/L	mmol/L
Serum potassium (K⁺) at time of testing? Serum potassium should be >2 mmol/L	mmol/L	mmol/L
Serum phosphate (PO₄³⁻) at time of testing? Serum phosphate should not be profoundly elevated (>3.0mmol/L) or lowered (<0.5mmol/L) from normal.	mmol/L	mmol/L
Serum magnesium (Mg²⁺) at time of testing? Serum magnesium should not be profoundly elevated (>3.0mmol/L) or lowered (<0.5mmol/L) from normal.	mmol/L	mmol/L
Blood glucose at time of testing? Blood glucose should be between 3.0-20.0 mmol/L and should be tested prior to each test.	mmol/L	mmol/L
If there is any clinical reason to expect endocrine disturbances hormonal assays should be undertaken.	Hormone level (if measured):	Hormone level (if measured):
Is the coma or apnoea due to a metabolic or endocrine disorder? (To diagnose death using neurological criteria, ALL answers should be NO)	Dr One	Dr Two
	Dr One	Dr Two

Guidance

It remains the duty of the two doctors carrying out the testing to be satisfied that the only explanation for the respiratory failure is due to the irreversible cessation of brain-stem function. A train of four examination, using a peripheral nerve stimulator, may be required. See above for 'Red Flag' patient groups.

	1 st Test		2 nd Test	
	Dr One	Dr Two	Dr One	Dr Two
Is the apnoea due to neuromuscular blocking agents, other drugs or a non brain-stem cause (e.g. cervical injury, any neuromuscular weakness)?				

Form for the Diagnosis of Death using Neurological Criteria in Children >2 months to <18 years {full guidance version}

Patient Name:

NHS Number:

Tests for Absence of Brain-Stem Function				
<p>Guidance: A complete set of tests should be performed on each occasion, i.e. a total of two sets of tests will be performed. Doctor One may perform the tests while Doctor Two observes; this would constitute the first set. Roles may be reversed for the second set. The tests, in particular the apnoea test, are therefore performed only twice in total.</p>				
	1 st Test		2 nd Test	
<p><u>To diagnose death using neurological criteria, ALL answers should be NO</u></p>	Dr One Examining	Dr Two Observing	Dr One Observing	Dr Two Examining
<p>Do the pupils react to light? The pupils are fixed and do not respond to sharp changes in the intensity of incident light. Cranial nerves II, III.</p>				
<p>Is there any eyelid movement when each cornea is touched in turn? Corneal reflex - Cranial nerves V, VII. The use of sterile gauze is recommended.</p>	Yes / No	Yes / No	Yes / No	Yes / No
<p>Is there any eye movement seen during or following the slow injection of at least 20 - 50mls ice cold water over 1 minute into each ear with the head at 30°? Each ear drum should be clearly visualised before the test. Vestibulo-ocular reflex - Cranial nerves III, VI, VIII.</p>	Yes / No	Yes / No	Yes / No	Yes / No
<p>Is the gag reflex present? Use a spatula or Yankauer sucker and laryngoscope to stimulate the posterior pharynx. Cranial Nerves IX, X.</p>				
<p>Is the cough reflex response present when a suction catheter is passed down the trachea to the carina? Cranial Nerves IX, X</p>				
<p>Is there any motor response in a cranial nerve or somatic distribution when supraorbital pressure is applied? Cranial Nerves V, VII. Reflex limb and trunk movements (spinal reflexes) can be present.</p>				

Tests for Absence of Brain-Stem Function
<p>Preparation for the Apnoea Test</p> <ul style="list-style-type: none"> Oxygenation and cardiovascular stability should be maintained through each apnoea test. Pre-oxygenate FiO₂ 1.0. Allow PaCO₂ to rise to at least 6.0 kPa by reducing the minute ventilation prior to commencing the apnoea test. End tidal carbon dioxide can be used to guide the starting of each apnoea test but should not replace the pre and post arterial PaCO₂. Cardiac pulsation may be sufficient to trigger supportive breaths if the patient remains connected to the mechanical ventilator and on a spontaneous breathing mode. Performing the apnoea test whilst remaining on mechanical ventilation is not recommended.

Form for the Diagnosis of Death using Neurological Criteria in Children >2 months to <18 years {full guidance version}

Patient Name:

NHS Number:

Guidance:

It is important to maintain oxygenation above 85% and cardiovascular stability.

Recommended methods:

- CPAP circuit (e.g. Mapleson C or Ayres T piece), especially if oxygenation is a problem, or
- Disconnect the patient from the ventilator and administer oxygen via a catheter in the trachea at a rate of 2- 6L/minute. Ensure oxygen catheter does not occlude ETT.
- Considerable atelectasis develops in the apnoeic period. At the conclusion of the apnoea test, manual recruitment manoeuvres should be carried out before resuming mechanical ventilation.

	1st Test		2nd Test	
Arterial Blood Gas PRE-apnoea test: Confirm PaCO ₂ is at least 6.0 kPa but not substantially greater. In patients with chronic CO ₂ retention, or those who have received intravenous bicarbonate, it recommended that PaCO ₂ is allowed to rise to above 6.5 kPa.	1 st Test Starting PaCO ₂ : <div style="text-align: right;">kPa</div> Should be ≥6.0 kPa		2 nd Test Starting PaCO ₂ : <div style="text-align: right;">kPa</div> Should be ≥6.0 kPa	
PRE-Arterial Blood Gas pH: Confirm pH < 7.4	pH= Should be < 7.4		pH= Should be < 7.4	
Start time: Time when apnoea test was commenced.	hr : min (24 hour clock)		hr : min (24 hour clock)	
Arterial Blood Gas POST apnoea test: Ensure the PaCO ₂ has increased by greater than 0.5 kPa.	1 st Test Stopping PaCO ₂ : <div style="text-align: right;">kPa</div> Should have increased by >0.5 kPa		2 nd Test Stopping PaCO ₂ : <div style="text-align: right;">kPa</div> Should have increased by >0.5 kPa	
Stop time: Time when apnoea test was ceased.	hr : min (24 hour clock)		hr : min (24 hour clock)	
	<i>Perform lung recruitment</i>		<i>Perform lung recruitment</i>	
Was there any spontaneous respiration during a minimum of 5 (five) minutes continuous observation following disconnection from the ventilator? (To diagnose death using neurological criteria, ALL answers should be NO)	Dr One	Dr Two	Dr One	Dr Two

Form for the Diagnosis of Death using Neurological Criteria in Children >2 months to <18 years {full guidance version}

Patient Name:

NHS Number:

Ancillary Investigations Used to Confirm the Diagnosis

Guidance:

Ancillary investigations are **NOT** required for the diagnosis and confirmation of death using neurological criteria.

They may be useful however, where neurological examination is not possible (e.g. extensive facio-maxillary injuries, residual sedation and some cases of paediatric hypoxic brain injury), where a primary metabolic or pharmacological derangement cannot be ruled out or in cases of high cervical cord injury, or where spontaneous or reflex movements in the patient generate uncertainty over the diagnosis. In such cases a confirmatory test may reduce any element of uncertainty and possibly foreshorten any period of observation prior to formal testing of brain-stem reflexes. Any ancillary or confirmatory investigation should be considered **ADDITIONAL** to the fullest clinical testing and examination (as outlined above) carried out to the best of the two doctors capabilities in the given circumstances.

The utility of any additional investigation is for the two testing doctors to decide and they should seek further professional opinion from other specialities and other expert centres, where appropriate. Some possible ancillary investigations are:

- Neurophysiological demonstration of loss of bioelectrical activity in the brain (EEG, evoked potentials).
- Radiological demonstration of absent cerebral blood flow or brain tissue perfusion (CT angiography, 4 vessel angiography, transcranial doppler).

The interpretation of ancillary investigations is complex and their availability usually restricted to neurological centres.

Helpful references on ancillary testing

1. Wijdicks (2001) "The Diagnosis of Brain Death" *NEJM* 344:1215-21.
2. Young & Lee (2004) "A critique of Ancillary Tests for Brain Death." *Neurocritical Care*; 1:499-508.
3. Heran, Heran & Shemie (2008) "A review of ancillary tests in evaluating brain death." *Can J Neurol Sci*; 35:409-19.

Is there a need for any ancillary investigations?	Dr One	Dr Two
---	---------------	---------------

If yes please outline the results of these investigations:

Form for the Diagnosis of Death using Neurological Criteria in Children >2 months to <18 years {full guidance version}

Patient Name:

NHS Number:

Completion of Diagnosis				
	Test 1		Test 2	
Are you satisfied that death has been confirmed following the irreversible cessation of brain-stem function?	Dr One	Dr Two	Dr One	Dr Two
Legal time of death is when the 1 st Test indicates death due to the absence of brain-stem reflexes.	Date: Time: Dr One signature		Date: Time: Dr One signature	
Death is confirmed following the 2 nd Test.	Dr Two signature		Dr Two signature	

Appendix 1 Paediatric Physiological Parameters⁵

Age	Heart Rate	Systolic BP	Mean Arterial BP
3 months	110 - 170	65 - 105	44 - 79
6 months	110 - 160	65 - 105	44 - 79
12 months	110 - 160	70 - 105	46 - 79
18 months	100 - 155	70 - 105	46 - 79
2 years	100 - 150	70 - 110	49 - 84
3 years	90 - 140	70 - 110	52 - 87
4 years	80 - 135	70 - 110	54 - 89
5 years	80 - 135	80 - 120	60 - 94
6 years	80 - 130	80 - 120	62 - 96
7 years	80 - 130	80 - 120	63 - 97
8 years	70 - 120	80 - 120	64 - 98
9 years	70 - 120	80 - 120	64 - 99
10 years	70 - 120	80 - 120	65 - 100
11 years	70 - 120	80 - 120	66 - 100
12 years	65 - 115	90 - 140	69 - 107
14 years	60 - 110	90 - 140	70 - 108
14 - 18 years	60 - 110	90 - 140	73 - 111

References

1. Academy of Medical Royal Colleges (2008) "A Code of Practice for the Diagnosis and Confirmation of Death" <http://www.aomrc.org.uk>
2. GMC (2010) "Treatment and care towards the end of life." www.gmc-uk.org/guidance/ethical_guidance/end_of_life_care.asp
3. NICE (2011, 2016) "Organ Donation for Transplantation" <http://guidance.nice.org.uk/CG135>
4. Paediatric Intensive Care Society (2014) "PICS Organ Donation Standards" <http://picsociety.uk/resources/>
5. Advanced Paediatric Life Support A Practical Approach to Emergencies (2016)
6. Royal College of Paediatrics and Child Health (2015) "The diagnosis of death by neurological criteria in infants less than two months old" www.rcpch.ac.uk

Form for the Diagnosis of Death using Neurological Criteria in Children >2 months to <18 years {full guidance version}

Patient Name:

NHS Number:

Form authorship and feedback

This form was written by Dr Dale Gardiner, Nottingham, Dr Alex Manara, Bristol and Dr Kay Hawkins, Manchester, Dr James Fraser, Bristol, Dr Margrid Shindler, Bristol and Andrea Macarthur, Manchester. Angie Scales, NHS Blood and Transplant. Comments should be directed to kay.hawkins@cmft.nhs.uk



Attach Arterial Blood Gases

Additional NOTES